

International Journal of Modern Physics A
 © World Scientific Publishing Company

Mixed QCD and Weak Corrections to $t\bar{t}$ Production by Quark-Antiquark Annihilation

W. Bernreuther ^{a,*}, M. Fűcker ^{a,*}, Z. G. Si ^{b,}

^a*Institut für Theoretische Physik, RWTH Aachen, 52056 Aachen, Germany*

^b*Department of Physics, Shandong University, Jinan, Shandong 250100, China*

We report on our computation of the mixed QCD and weak corrections to $q\bar{q} \rightarrow t\bar{t}$ including $t, \bar{t}$ spin effects, in particular on single top spin asymmetries.

Top quark physics will play an important role in testing the Standard Model (SM) and searching for new physics beyond the SM, once large $t\bar{t}$ data samples are available at the Tevatron and at the LHC in the future. For this aim, the SM predictions should be known as precisely as possible. Specifically weak interaction contributions to $t\bar{t}$ production should be taken into account. Although they are nominally subdominant with respect to the QCD contributions they can become important at large $t\bar{t}$ invariant mass due to large Sudakov logarithms (for reviews and references, see e.g. ^{1,2}).

SM weak interaction effects in hadronic production of heavy quark pairs were considered previously. The parity-even and parity-odd order $\alpha_s^2\alpha$ vertex corrections were determined in ³ and in ⁴, respectively (see also ⁵). In ref. ⁴ also parity-violating non-SM effects were analysed. The box contributions to $q\bar{q} \rightarrow t\bar{t}$ were not taken into account in these papers. In the following we report on the results of the mixed QCD and weak radiative corrections of order $\alpha_s^2\alpha$ to the (differential) cross section of $t\bar{t}$ production by $q\bar{q}$ annihilation, keeping the full information on the spin state of the $t\bar{t}$ system. These are necessary ingredients for definite SM predictions, in particular of parity-violating observables associated with the spin of the (anti)top quark.

Top quark pair production both at the Tevatron and at the LHC is dominated by the QCD contributions to $q\bar{q} \rightarrow t\bar{t}$ and $gg \rightarrow t\bar{t}$. Their cross section including the full dependence on the $t, \bar{t}$ spins are known to order (α_s^3) ^{6,7}. Due to color conservation, the leading contributions involving electroweak interactions are the order α^2 Born terms for $q\bar{q} \rightarrow t\bar{t}$ and the mixed contributions of order $\alpha_s^2\alpha$. For the $q + \bar{q} \rightarrow t + \bar{t}$, and $q + \bar{q} \rightarrow t + \bar{t} + g$ processes, the inclusive spin-summed cross sections may be written, to NLO in the SM couplings, in the form

$$\sigma_{q\bar{q}} = \sigma_{q\bar{q}}^{(0)QCD} + \delta\sigma_{q\bar{q}}^{QCD} + \delta\sigma_{q\bar{q}}^W, \quad (1)$$

where the first and second term are the LO (order α_s^2) and NLO (order α_s^3) QCD contributions ^{8,9,10}. The third term is generated by the electroweak contributions,

2 *W. Bernreuther, M. Fückler, Z. G. Si*

and can be decomposed as follows:

$$\delta\sigma_{q\bar{q}}^W(\hat{s}, m_t^2) = \frac{4\pi\alpha}{m_t^2} [\alpha f_{q\bar{q}}^{(0)}(\eta) + \alpha_s^2 f_{q\bar{q}}^{(1)}(\eta)], \quad \eta = \frac{\hat{s}}{4m_t^2} - 1, \quad (2)$$




Fig. 1. (a) Dimensionless scaling functions $f_{q\bar{q}}^{(0)}(\eta)$ (dashed), $f_{q\bar{q}}^{(1)}(\eta)$ (solid) that determine the parton cross section (2) for $q = d$ type quarks. The dash-dotted and dotted lines correspond to the respective functions for $q = u$ type quarks. (b) Contributions of the LO (solid) and NLO QCD (dashed) contributions (taken from ⁷) and of the mixed $\alpha_s^2\alpha$ contributions (dotted and dash-dotted line refers to initial d-type and u-type quarks, respectively) to the cross section (1) in units of $1/m_t^2$.

where $\hat{s}$ is the $q\bar{q}$ c.m. energy squared. In our calculations ¹¹ we used $m_t = 178$ GeV, $m_Z = 91.188$ GeV, $\sin^2\theta_W = 0.231$, and $m_H = 114$ GeV for the mass of the SM Higgs boson. In Fig. 1(a) the functions $f_{q\bar{q}}^{(i)}$ are displayed as functions of η for annihilation of initial massless partons $q\bar{q}$ of the first and second generation with weak isospin $\pm 1/2$. As expected the $\alpha_s^2\alpha$ corrections are significantly larger than the the lowest order photon and Z boson exchange contributions. The correction (2) to the $q\bar{q}$ cross section has recently been computed also by ¹². We have compared our results and find excellent numerical agreement. The contribution of the corrections shown in Fig. 1(a) to the $t\bar{t}$ cross section at the Tevatron is very small. This is mainly due to the fact that the order $\alpha_s^2\alpha$ corrections change sign for larger η . The significance of the contributions can be enhanced by suitable cuts, e.g. in the $t\bar{t}$ invariant mass. In Fig. 1(b) the order α_s^2 , α_s^3 , and the $\alpha_s^2\alpha$ contributions to the cross section (1) are shown as functions of η . In these plots $\alpha_s(m_t) = 0.095$ and $\alpha(m_Z) = 0.008$ was chosen. One sees that for $\eta \gtrsim 1$ the mixed corrections become of the same size or larger in magnitude than the NLO QCD contributions, and at $\eta \sim 10$ the $\alpha_s^2\alpha$ contributions are already about 15 percent of the LO QCD cross section. These regions can be investigated by studying the distribution of the $t\bar{t}$ invariant mass $M_{t\bar{t}}$.

Next we consider spin asymmetries. Denoting the top spin operator by $\mathbf{S}_t$ and its projection onto an arbitrary unit axis $\hat{\mathbf{a}}$ by $\mathbf{S}_t \cdot \hat{\mathbf{a}}$ we can express its unnormalized partonic expectation value, which we denote by double brackets, in terms of the

difference between the “spin up” and “spin down” cross sections:

$$2\langle\langle\mathbf{S}_t \cdot \hat{\mathbf{a}}\rangle\rangle_i = \sigma_i(\uparrow) - \sigma_i(\downarrow). \quad (3)$$

Here $i = q\bar{q}$ and the arrows refer to the spin state of the top quark with respect to $\hat{\mathbf{a}}$. An analogous formula holds for the antitop quark. It is these expressions that enter the predictions for (anti)proton collisions.

There are two types of single spin asymmetries (3): parity-even, T-odd asymmetries and parity-violating, T-even ones. The asymmetry associated with the projection $\mathbf{S}_t$ onto the normal of the q, t scattering plane belongs to the first class. It is induced by the absorptive part of the amplitude which receives a QCD contribution at NLO ^{13,14}. The weak contribution is even smaller than the one from QCD which is of the order of a few percent. For the sake of brevity we do not display it here. The P-odd, T-even single spin asymmetries correspond to projections of




Fig. 2. Scaling functions $h_{q\bar{q}}^{(0,a)}(\eta)$ (dashed), $h_{q\bar{q}}^{(1,a)}(\eta)$ (solid) that determine the expectation value (4) for (a) beam and (b) helicity axis in the case of $q = d$ type quarks. The dash-dotted and dotted lines correspond to the respective functions for $q = u$ type quarks.

the top spin onto a polar vector, in particular onto an axis that lies in the scattering plane. Needless to say, they cannot be generated within QCD; the leading SM contributions to these asymmetries are the parity-violating pieces. We consider top spin projections onto the beam axis (which is relevant for the Tevatron), and onto the helicity axis (of relevance for the LHC). Naively, one might define these axes in the c.m. frame of the initial partons. However, the observables $\mathbf{S}_t \cdot \hat{\mathbf{a}}$ are then not collinear-safe. (The problem shows up once second-order QCD corrections are taken into account.) A convenient frame with respect to which collinear-safe spin projections can be defined is the $t\bar{t}$ zero-momentum-frame (ZMF) ⁷. With respect to this frame we define the axes as $\hat{\mathbf{a}} = \hat{\mathbf{p}}$ (beam basis) and $\hat{\mathbf{a}} = \hat{\mathbf{k}}$ (helicity axes) where $\hat{\mathbf{p}}$ is the direction of flight of one of the colliding hadrons, and $\hat{\mathbf{k}}$ denotes the direction of flight of the top quark. The unnormalized expectation values of $\mathbf{S}_t \cdot \hat{\mathbf{a}}$ are again conveniently expressed by scaling functions

$$\langle\langle 2\mathbf{S}_t \cdot \hat{\mathbf{a}} \rangle\rangle_{q\bar{q}} = \frac{4\pi\alpha}{m_t^2} [\alpha h_{q\bar{q}}^{(0,a)}(\eta) + \alpha_s^2 h_{q\bar{q}}^{(1,a)}(\eta)]. \quad (4)$$

4 *W. Bernreuther, M. Fückler, Z. G. Si*

The results for the scaling functions corresponding to beam and helicity axes are shown in Figs. 2(a) and (b). For the beam axes, and for weak isospin $I_W = -1/2$ quarks the $\alpha_s^2\alpha$ corrections are significantly larger than the LO values. This is in contrast to the helicity basis where the LO and NLO terms shown in Fig. 2(b) are of the same order of magnitude. Moreover, in this basis the LO and NLO terms cancel each other to a large extent for $I_W = -1/2$ quarks in the initial state. Because $t\bar{t}$ production at the Tevatron occurs predominantly by $q\bar{q}$ annihilation, these results imply that, when it comes to searching for SM-induced parity-violating t or $\bar{t}$ spin effects, the reference axes of choice should be $\hat{\mathbf{p}}$. The SM effect is quite small; for suitably chosen $M_{t\bar{t}}$ mass bins one gets asymmetries of the order of 2 percent. This leaves a large margin in the search for new physics contributions.

To summarize, we have investigated the mixed QCD and weak interaction contributions to the cross section of $q\bar{q} \rightarrow t\bar{t}$ and computed the parity-violating single (anti)top quark spin effects induced by these interactions (for details, see ¹¹). The corresponding investigation of $gg \rightarrow t\bar{t}$, which is necessary for a comprehensive analysis of $pp, p\bar{p} \rightarrow t\bar{t} X$, is in progress. The parity-violating observables associated with the spin of the (anti)top quark are very sensitive to non-SM parity-violating top quark interactions, and therefore they will become very important analysis tools once sufficiently large $t\bar{t}$ data samples are available at the Tevatron and at the LHC.

Acknowledgements

This work was supported by DFG SFB/TR9, by DFG-Graduiertenkolleg RWTH Aachen, and by NSFC and NCET of China.

References

1. M. Melles, Phys. Rept. **375** (2003) 219.
2. A. Denner, arXiv:hep-ph/0110155.
3. W. Beenakker, A. Denner, W. Hollik, R. Mertig, T. Sack and D. Wackeroth, Nucl. Phys. B **411** (1994) 343.
4. C. Kao and D. Wackeroth, Phys. Rev. D **61** (2000) 055009.
5. C. Kao, G. A. Ladinsky and C. P. Yuan, Int. J. Mod. Phys. A **12** (1997) 1341.
6. W. Bernreuther, A. Brandenburg, Z. G. Si and P. Uwer, Phys. Rev. Lett. **87** (2001) 242002.
7. W. Bernreuther, A. Brandenburg, Z. G. Si and P. Uwer, Nucl. Phys. B **690** (2004) 81.
8. P. Nason, S. Dawson and R. K. Ellis, Nucl. Phys. B **303** (1988) 607.
9. W. Beenakker, W. L. van Neerven, R. Meng, G. A. Schuler and J. Smith, Nucl. Phys. B **351** (1991) 507.
10. W. Bernreuther, A. Brandenburg and Z. G. Si, Phys. Lett. B **483** (2000) 99.
11. W. Bernreuther, M. Fückler and Z. G. Si, arXiv:hep-ph/0508091.
12. J. H. Kühn, A. Scharf and P. Uwer, arXiv:hep-ph/0508092.
13. W. Bernreuther, A. Brandenburg and P. Uwer, Phys. Lett. B **368** (1996) 153.
14. W. G. Dharmaratna and G. R. Goldstein, Phys. Rev. D **53** (1996) 1073.